


Amendments to the Claims:

The following is a complete list of claims indicating the changes incorporated by the present amendment and replacing all prior versions of the claims. Any claims canceled herein and all deletions made in claims that are not canceled herein are done so without prejudice to being re-instituted at a later date in this or a related application.

Listing of Claims:

- 1 1. (Currently Amended) A compound having the formula:



(I)

2 wherein:

3 L² and L⁴ are members independently selected from a bond, substituted or
4 unsubstituted alkylene, and substituted or unsubstituted heteroalkylene;
5 L³ is a member selected from a bond, substituted or unsubstituted alkylene,
6 substituted or unsubstituted heteroalkylene, -C(O)-, -C(O)NH-, and -S(O)_u-,
7 wherein u is 0, 1, or 2;
8 the dashed lines a and b are optionally a bond, wherein if R² is =O, =N-OR^{2A}, or
9 =CR^{2B}R^{2C}, then R¹ is absent, L² is a bond, and a is a bond attached directly
10 to R²;
11 R¹ is absent or a member selected from hydrogen, substituted or unsubstituted
12 alkyl, substituted or unsubstituted heteroalkyl, substituted or unsubstituted
13 cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or
14 unsubstituted aryl, and substituted or unsubstituted heteroaryl;
15 R² is a member selected from =O, =N-OR^{2A}, =CR^{2B}R^{2C}, hydrogen, -OR^{2D},
16 -C(O)R^{2D}, -C(O)NR^{2E}R^{2F}, -NR^{2E}R^{2F}, substituted or unsubstituted alkyl,

18 substituted or unsubstituted heteroalkyl, substituted or unsubstituted
19 cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or
20 unsubstituted aryl, and substituted or unsubstituted heteroaryl, wherein
21 R^{2A}, R^{2B}, R^{2C} and R^{2D} are members independently selected from hydrogen,
22 substituted or unsubstituted alkyl, substituted or unsubstituted
23 heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or
24 unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and
25 substituted or unsubstituted heteroaryl;
26 R^{2E} and R^{2F} are members independently selected from hydrogen, substituted
27 or unsubstituted alkyl, substituted or unsubstituted heteroalkyl,
28 substituted or unsubstituted aryl, substituted or unsubstituted
29 heterocycloalkyl, -S(O)_mR^{2E1} and -S(O)_mNR^{2E2}R^{2E3}, wherein
30 R^{2E} and R^{2F} are optionally joined to form a substituted or unsubstituted
31 ring with the nitrogen to which they are attached, and
32 R^{2E1}, R^{2E2}, and R^{2E3} are members independently selected from hydrogen,
33 substituted or unsubstituted alkyl, substituted or unsubstituted
34 heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or
35 unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and
36 substituted or unsubstituted heteroaryl, and
37 m is 0, 1, or 2,
38 wherein R² and R¹ are optionally joined to form a substituted or
39 unsubstituted ring;
40 R³ is a member selected from substituted or unsubstituted higher alkyl,
41 substituted or unsubstituted heteroalkyl, substituted or unsubstituted
42 cycloalkyl, substituted or unsubstituted heterocycloalkyl, substituted or
43 unsubstituted aryl, and substituted or unsubstituted heteroaryl, ~~-OR^{3A}, and-~~
44 ~~NR^{3B}R^{3C}, wherein~~
45 ~~R^{3A}, R^{3B}, and R^{3C} are independently members selected from substituted~~
46 ~~or unsubstituted higher alkyl, substituted or unsubstituted~~


47 heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or
48 unsubstituted heterocycloalkyl, substituted or unsubstituted aryl,
49 and substituted or unsubstituted heteroaryl, wherein R^{3B} and R^{3C}
50 are optionally joined to form a substituted or unsubstituted ring
51 with the nitrogen to which they are attached;

52 R⁴ is a member selected from substituted or unsubstituted alkyl, substituted
53 or unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl,
54 substituted or unsubstituted heterocycloalkyl, substituted or
55 unsubstituted aryl, substituted or unsubstituted heteroaryl, S(O)_tR^{4A},
56 S(O)_tNR^{4B}R^{4C}, C(O)R^{4A}, C(O)OR^{4A}, C(O)NR^{4B}R^{4C}, =NOR^{4D}, and
57 =NOR^{4E}NR^{4F}, wherein

58 R^{4A}, R^{4B}, R^{4C}, R^{4D}, R^{4E}, and R^{4F} are members independently selected
59 from substituted or unsubstituted alkyl, substituted or unsubstituted
60 heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or
61 unsubstituted heterocycloalkyl, substituted or unsubstituted aryl,
62 substituted or unsubstituted heteroaryl, and

63 t is 0, 1, or 2.

64 R⁴ has the formula:



66 wherein

67 R^{4G} is a member independently selected from hydrogen, halogen, -OH,
68 -COOH, -CF₃, -NH₂, -SH, substituted or unsubstituted alkyl,
69 substituted or unsubstituted heteroalkyl, substituted or
70 unsubstituted cycloalkyl, substituted or unsubstituted
71 heterocycloalkyl, substituted or unsubstituted aryl, and substituted
72 or unsubstituted heteroaryl;

73 A is a substituted or unsubstituted ring selected from substituted or
74 unsubstituted (C₃-C₇) cycloalkyl, substituted or unsubstituted 3-7
75 membered heterocycloalkyl, substituted or unsubstituted aryl, and
76 substituted or unsubstituted heteroaryl;
77 X is a member selected from a bond, -S(O)_v-, and -S(O)_vNR^{4I}-, wherein
78 R^{4I} is a member selected from hydrogen, substituted or
79 unsubstituted alkyl, and substituted or unsubstituted heteroalkyl,
80 and
81 v is 0, 1, or 2; and
82 w is an integer from 1 to 5.

1 2. (Original) The compound of claim 1, wherein
2 R² is a member selected from =O, =N-OR^{2A}, -OR^{2D}, -NR^{2E}R^{2F}, substituted or
3 unsubstituted (C₁-C₁₀) alkyl, substituted or unsubstituted 2-10 membered
4 heteroalkyl, substituted or unsubstituted (C₃-C₇) cycloalkyl, substituted or
5 unsubstituted 3-7 membered heterocycloalkyl, substituted or unsubstituted
6 aryl, and substituted or unsubstituted heteroaryl, wherein
7 R^{2A} and R^{2D} are members independently selected from hydrogen and
8 substituted or unsubstituted (C₁-C₁₀) alkyl, and
9 R^{2E} and R^{2F} are members independently selected from hydrogen and
10 substituted or unsubstituted (C₁-C₁₀) alkyl.

1 3. (Original) The compound of claim 1, wherein
2 R² is a member selected from =O, =N-OR^{2A}, and -OR^{2D}, wherein
3 R^{2A} and R^{2D} are members independently selected from hydrogen and
4 unsubstituted (C₁-C₅) alkyl.

1 4. (Original) The compound of claim 1, wherein R² is =O and the dashed
2 line b is a bond.


1 **5.** (Original) The compound of claim 1, wherein R¹ is absent or is a
2 member selected from hydrogen and substituted or unsubstituted alkyl.

1 **6.** (Original) The compound of claim 1, wherein R¹ is absent or is a
2 member selected from hydrogen, methyl, and -C≡C-CH₃.

1 **7.** (Original) The compound of claim 1, wherein R¹ is absent.

1 **8.** (Original) The compound of claim 1, wherein R³ is a member selected
2 from substituted or unsubstituted (C₁-C₁₀) alkyl, substituted or unsubstituted 2-10 membered
3 heteroalkyl, substituted or unsubstituted (C₃-C₇) cycloalkyl, substituted or unsubstituted 3-7
4 membered heterocycloalkyl, substituted or unsubstituted aryl, and substituted or unsubstituted
5 heteroaryl.

1 **9.** (Original) The compound of claim 1, wherein R³ has the formula:



(III)

3 wherein

4 q is an integer selected from 1 to 5; and

5 R^{3D} is a member independently selected from hydrogen, halogen, -OH, -COOH,
6 -CF₃, -NH₂, -SH, substituted or unsubstituted alkyl, substituted or
7 unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted
8 or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl,
9 substituted or unsubstituted heteroaryl, -NR^{3D1}R^{3D2}, -OR^{3D3},
10 -C(O)NR^{3D4}R^{3D5}, and -C(O)R^{3D6}, wherein

11 R^{3D1}, R^{3D2}, R^{3D3}, R^{3D4}, R^{3D5}, and R^{3D6} are members independently selected from
12 hydrogen, substituted or unsubstituted alkyl, substituted or unsubstituted
13 heteroalkyl, substituted or unsubstituted cycloalkyl, substituted or
14 unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and
15 substituted or unsubstituted heteroaryl, wherein


16 R^{3D1} and R^{3D2} are optionally joined to form a substituted or unsubstituted
17 ring with the nitrogen to which they are attached, and
18 R^{3D4} and R^{3D5} are optionally joined to form a substituted or unsubstituted
19 ring with the nitrogen to which they are attached.

1 **10.** (Original) The compound of claim 9, wherein
2 q is an integer selected from 1 to 3; and
3 R^{3D} is a member independently selected from hydrogen, substituted alkyl,
4 substituted or unsubstituted heteroalkyl, substituted cycloalkyl, substituted
5 or unsubstituted heterocycloalkyl, substituted aryl, and substituted or
6 unsubstituted heteroaryl.

1 **11.** (Original) The compound of claim 9, wherein
2 R^{3D} is a member independently selected from hydrogen, R^{3D7} -substituted (C_1 -
3 C_{10}) alkyl, R^{3D7} -substituted or unsubstituted 2-10 membered heteroalkyl,
4 R^{3D7} -substituted or unsubstituted (C_3 - C_8) cycloalkyl, R^{3D7} -substituted or
5 unsubstituted 3-8 membered heterocycloalkyl, R^{3D8} -substituted or
6 unsubstituted aryl, R^{3D8} -substituted or unsubstituted heteroaryl, $-NR^{3D1}R^{3D2}$,
7 $-OR^{3D3}$, $-C(O)NR^{3D4}R^{3D5}$, and $-C(O)R^{3D6}$, wherein
8 R^{3D1} , R^{3D2} , R^{3D3} , R^{3D4} , R^{3D5} , and R^{3D6} are members independently selected
9 from hydrogen, R^{3D7} -substituted or unsubstituted alkyl, R^{3D7} -substituted
10 or unsubstituted heteroalkyl, R^{3D7} -substituted or unsubstituted
11 cycloalkyl, R^{3D7} -substituted or unsubstituted heterocycloalkyl, R^{3D8} -
12 substituted or unsubstituted aryl, and R^{3D8} -substituted or unsubstituted
13 heteroaryl,
14 wherein R^{3D1} and R^{3D2} are optionally joined with the nitrogen to which
15 they are attached to form a R^{3D7} -substituted or unsubstituted
16 heterocycloalkyl, or R^{3D8} -substituted or unsubstituted heteroaryl, and

17 wherein R^{3D4} and R^{3D5} are optionally joined with the nitrogen to which
18 they are attached to form a R^{3D7}-substituted or unsubstituted
19 heterocycloalkyl, or R^{3D8}-substituted or unsubstituted heteroaryl,
20 wherein
21 R^{3D7} is a member selected from halogen, oxo, -OH, -COOH, -CF₃, -NH₂,
22 -SH, R^{3D9}-substituted or unsubstituted (C₁-C₁₀) alkyl, R^{3D9}-substituted or
23 unsubstituted 2-10 membered heteroalkyl, R^{3D9}-substituted or
24 unsubstituted (C₃-C₈) cycloalkyl, R^{3D9}-substituted or unsubstituted 3-8
25 membered heterocycloalkyl, R^{3D10}-substituted or unsubstituted aryl, and
26 R^{3D10}-substituted or unsubstituted heteroaryl, and
27 R^{3D8} is a member selected from halogen, -OH, -COOH, -CF₃, -NH₂, -SH,
28 R^{3D9}-substituted or unsubstituted (C₁-C₁₀) alkyl, R^{3D9}-substituted or
29 unsubstituted 2-10 membered heteroalkyl, R^{3D9}-substituted or
30 unsubstituted (C₃-C₈) cycloalkyl, R^{3D9}-substituted or unsubstituted 3-8
31 membered heterocycloalkyl, R^{3D10}-substituted or unsubstituted aryl, and
32 R^{3D10}-substituted or unsubstituted heteroaryl,
33 R^{3D9} is a member selected from halogen, oxo, -OH, -COOH, -CF₃, -NH₂,
34 -SH, unsubstituted (C₁-C₁₀) alkyl, unsubstituted 2-10 membered
35 heteroalkyl, unsubstituted (C₃-C₈) cycloalkyl, unsubstituted 3-8
36 membered heterocycloalkyl, unsubstituted aryl, and unsubstituted
37 heteroaryl, and
38 R^{3D10} is a member selected from halogen, -OH, -COOH, -CF₃, -NH₂, -SH,
39 unsubstituted (C₁-C₁₀) alkyl, unsubstituted 2-10 membered heteroalkyl,
40 unsubstituted (C₃-C₈) cycloalkyl, unsubstituted 3-8 membered
41 heterocycloalkyl, unsubstituted aryl, and unsubstituted heteroaryl.

1 12. (Original) The compound of claim 11, wherein R³ has the formula:



3 wherein

4 R^{3D} is a member selected from hydrogen, R^{3D7}-substituted (C₁-C₅) alkyl, R^{3D7}-
5 substituted or unsubstituted 2-5 membered heteroalkyl, R^{3D7}-substituted (C₅-
6 C₇)cycloalkyl, R^{3D7}-substituted or unsubstituted 5-7 membered
7 heterocycloalkyl, R^{3D8}-substituted aryl, R^{3D8}-substituted or unsubstituted
8 heteroaryl, -NR^{3D1}R^{3D2}, -OR^{3D3}, -C(O)NR^{3D4}R^{3D5}, and -C(O)R^{3D6}.

1 13. (Original) The compound of claim 12, wherein R^{3D} is a member selected
2 from -NR^{3D1}R^{3D2}, -OR^{3D3}, -C(O)NR^{3D4}R^{3D5}, and R^{3D7}-substituted or unsubstituted heteroaryl
3 comprising a ring nitrogen, wherein

4 R^{3D1} and R^{3D2} are members independently selected from hydrogen, R^{3D7}-
5 substituted alkyl, R^{3D7}-substituted or unsubstituted heteroalkyl, R^{3D7}-
6 substituted or unsubstituted heterocycloalkyl, and R^{3D8}-substituted or
7 unsubstituted heteroaryl,
8 wherein R^{3D1} and R^{3D2} are optionally joined with the nitrogen to which they
9 are attached to form a R^{3D7}-substituted or unsubstituted heterocycloalkyl,
10 or R^{3D8}-substituted or unsubstituted heteroaryl, wherein said ring
11 optionally comprises an additional ring heteroatom; and
12 R^{3D3}, R^{3D4} and R^{3D5} are members independently selected from
13 hydrogen,
14 R^{3D7}-substituted or unsubstituted heteroalkyl comprising a nitrogen
15 heteroatom,
16 R^{3D7}-substituted or unsubstituted heterocycloalkyl comprising a ring
17 nitrogen,
18 R^{3D8}-substituted or unsubstituted heteroaryl comprising a ring nitrogen, and
19 alkyl substituted with a R^{3D9}-substituted or unsubstituted heteroalkyl
20 comprising a nitrogen heteroatom, R^{3D9}-substituted or unsubstituted
21 heterocycloalkyl comprising a ring nitrogen, or R^{3D10}-substituted or
22 unsubstituted heteroaryl comprising a ring nitrogen,

23 wherein R^{3D4} and R^{3D5} are optionally joined with the nitrogen to which they
24 are attached to form a R^{3D7}-substituted or unsubstituted heterocycloalkyl,
25 or R^{3D8}-substituted or unsubstituted heteroaryl, wherein said ring
26 optionally comprises a heteroatom.

1 **14.** (Original) The compound of claim 13, wherein
2 R^{3D1} and R^{3D2}, and R^{3D4} and R^{3D5} are optionally joined with the nitrogen to
3 which they are attached to form a R^{3D7}-substituted or unsubstituted
4 heterocycloalkyl comprising an additional heteroatom, or R^{3D8}-substituted or
5 unsubstituted heteroaryl comprising an additional heteroatom.

1 **15.** (Original) The compound of claim 14, wherein R^{3D1} and R^{3D2}, and R^{3D4}
2 and R^{3D5} are optionally joined with the nitrogen to which they are attached to form a R^{3D8}-
3 substituted or unsubstituted oxazolyl, imidazolyl, thiazolyl, isooxazolyl, pyrazolyl, isothiazolyl,
4 purinyl, pyradizinyl, pyrimidinyl, pyrazinyl, or quinoxalinyl.

1 **16.-18.** (Canceled)

1 **19.** (Currently Amended) The compound of claim 1 [[18]], wherein
2 R^{4G} is a member independently selected from hydrogen, halogen, -OH, -COOH,
3 -CF₃, -NH₂, -SH, R^{4G1}-substituted or unsubstituted alkyl, R^{4G1}-substituted or
4 unsubstituted heteroalkyl, R^{4G1}-substituted or unsubstituted cycloalkyl, R^{4G1}-
5 substituted or unsubstituted heterocycloalkyl, R^{4G2}-substituted or
6 unsubstituted aryl, and R^{4G2}-substituted or unsubstituted heteroaryl, wherein
7 R^{4G1} is a member selected from halogen, oxo, -OH, -COOH, -CF₃, -NH₂,
8 -SH, R^{4G3}-substituted or unsubstituted (C₁-C₁₀) alkyl, R^{4G3}-substituted or
9 unsubstituted 2-10 membered heteroalkyl, R^{4G3}-substituted or
10 unsubstituted (C₃-C₈) cycloalkyl, R^{4G3}-substituted or unsubstituted 3-8
11 membered heterocycloalkyl, R^{4G4}-substituted or unsubstituted aryl, and
12 R^{4G4}-substituted or unsubstituted heteroaryl, and

13 R^{4G2} is a member selected from halogen, -OH, -COOH, -CF₃, -NH₂, -SH,
14 R^{4G3} -substituted or unsubstituted (C₁-C₁₀) alkyl, R^{4G3} -substituted or
15 unsubstituted 2-10 membered heteroalkyl, R^{4G3} -substituted or
16 unsubstituted (C₃-C₈) cycloalkyl, R^{4G3} -substituted or unsubstituted 3-8
17 membered heterocycloalkyl, R^{4G4} -substituted or unsubstituted aryl, and
18 R^{4G4} -substituted or unsubstituted heteroaryl,
19 R^{4G3} is a member selected from halogen, oxo, -OH, -COOH, -CF₃, -NH₂,
20 -SH, unsubstituted (C₁-C₁₀) alkyl, unsubstituted 2-10 membered
21 heteroalkyl, unsubstituted (C₃-C₈) cycloalkyl, unsubstituted 3-8
22 membered heterocycloalkyl, unsubstituted aryl, and unsubstituted
23 heteroaryl, and
24 R^{4G4} is a member selected from halogen, -OH, -COOH, -CF₃, -NH₂, -SH,
25 unsubstituted (C₁-C₁₀) alkyl, unsubstituted 2-10 membered heteroalkyl,
26 unsubstituted (C₃-C₈) cycloalkyl, unsubstituted 3-8 membered
27 heterocycloalkyl, unsubstituted aryl, and unsubstituted heteroaryl.

1 **20.** (Original) The compound of claim 19, wherein A is a member selected
2 from phenyl, pyrazolyl, furanyl, imidazolyl, isoxazolyl, oxadiazolyl, oxazolyl, pyrrolyl,
3 pyridyl, pyrazyl, pyrimidyl, pyridazinyl, thiazolyl, isothioazolyl, triazolyl, thienyl, triazinyl,
4 thiadiazolyl, dioxolanyl, dioxanyl, trioxanyl, tetrahydrothienyl, tetrahydrofuranyl,
5 tetrahydrothiophenyl, tetrahydropyranyl, tetrahydrothiopyranyl, pyrrolidinyl, morpholino,
6 piperidinyl, and piperazinyl.

1 **21.** (Currently Amended) The compound of claim 1 [[18]], wherein
2 R^{4G} is selected from hydrogen, substituted (C₁-C₅) alkyl, substituted or
3 unsubstituted 2-5 membered heteroalkyl, substituted (C₅-C₇)cycloalkyl,
4 substituted or unsubstituted heterocycloalkyl, substituted aryl, and
5 substituted or unsubstituted heteroaryl;

6 A is a substituted or unsubstituted ring selected from substituted or unsubstituted
7 3-7 membered heterocycloalkyl, substituted or unsubstituted aryl, and
8 substituted or unsubstituted heteroaryl; and
9 R⁴¹ is hydrogen.

1 22. (Currently Amended) The compound of claim 1 [[18]], wherein R^{4G} is a
2 branched or unbranched (C₁-C₁₀)alkyl.

1 23. (Currently Amended) The compound of claim 1 [[18]], wherein X is
2 -S(O)₂-.

1 24. (Currently Amended) The compound of claim 1, wherein [.-]
2 L², L³ and L⁴ are members independently selected from a bond and
3 unsubstituted (C₁-C₅) alkylene.

1 25. (Currently Amended) The compound of claim 1 wherein
2 the dashed line b is a bond;
3 R² is =O;
4 R³ is substituted or unsubstituted benzyl;
5 **R⁴ has the formula:**



7 **wherein**

8 R^{4G} is a member selected from substituted or unsubstituted alkyl,
9 substituted or unsubstituted heteroalkyl, substituted or
10 unsubstituted cycloalkyl, substituted or unsubstituted
11 heterocycloalkyl, substituted or unsubstituted aryl, and
12 substituted or unsubstituted heteroaryl,
13 **A is a substituted or unsubstituted ring selected from substituted or**
14 **unsubstituted (C₃-C₇) cycloalkyl, substituted or unsubstituted**

15 ~~3-7 membered heterocycloalkyl, substituted or unsubstituted~~
16 ~~aryl, and substituted or unsubstituted heteroaryl, and~~

17 X is -S(O)₂-;

18 W is 1;

19 L³ is a bond; and

20 L⁴ is a bond.


1 26. (Original) The compound of claim 1 having the formula:




1 27. (Original) The compound of claim 1 having the formula:



1 28. (Original) The compound of claim 1 having the formula:



1 29. (Currently Amended) The compound of claim 1 having the formula:



3 wherein

4 R^{4G} is a member selected from substituted or unsubstituted alkyl, substituted or
5 unsubstituted heteroalkyl, substituted or unsubstituted cycloalkyl, substituted
6 or unsubstituted heterocycloalkyl, substituted or unsubstituted aryl, and
7 substituted or unsubstituted heteroaryl;

8 **A is a substituted or unsubstituted ring selected from substituted or**
9 **unsubstituted (C₃-C₇) cycloalkyl, substituted or unsubstituted 3-7**
10 **membered heterocycloalkyl, substituted or unsubstituted aryl, and**
11 **substituted or unsubstituted heteroaryl;** and

12 X is a member selected from a bond, -S(O)₂-, and -S(O)₂NR^{4I}, wherein
13 R^{4I} is a member selected from hydrogen, substituted or unsubstituted alkyl,
14 and substituted or unsubstituted heteroalkyl.


1 30. (Original) A method of treating a disorder or condition through
2 modulating a glucocorticoid receptor, the method comprising administering to a subject in need
3 of such treatment, an effective amount of the compound of claim 1.

1 31. (Original) A method of treating a disorder or condition through
2 antagonizing a glucocorticoid receptor, the method comprising administering to a subject in
3 need of such treatment, an effective amount of the compound of claim 1.

1 32. (Original) A method of modulating a glucocorticoid receptor including
2 the steps of contacting a glucocorticoid receptor with the compound of claim 1 and detecting a
3 change in the activity of the glucocorticoid receptor.

1 33. (Original) A pharmaceutical composition comprising a pharmaceutically
2 acceptable excipient and the compound of claim 1.





1 34. (New) The compound of claim 1 having the formula:















3 wherein







4 R^{4A} is selected from the group consisting of cycloalkyl, heterocycloalkyl, aryl, and
5 heteroaryl.







1 35. (New) The compound of claim 1, selected from the group consisting of:























(XLVII)




(XLVIII)




(XLIX)









(L)














(LI)








(LII)














(LXXIV)



(LXXV)



(LXXVI)



(LXXVII)

